

AD A032670

AFOSR - TR - 76 - 1158

A CLASS OF CONTROLLABLE NONLINEAR SYSTEMS*

by

Kuang-Chung Wei
Division of Engineering
and
Lefschetz Center for Dynamical Systems
Brown University
Providence, RI 02912

Abstract

Sufficient conditions for global and local controllability of the nonlinear system

$$\frac{d}{dt} x(t) = A(t, x(t), u(t))x(t) + B(t)u(t) + f(t, x(t), u(t))$$

and its perturbed system are given. These conditions extend some previous results through the removal of certain boundedness conditions involving the functions (A, f) and their partial derivatives.

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----

Approved for public release;
distribution unlimited.

*This research was supported in part by the Air Force Office of Scientific Research under Grant AFOSR-75-2793.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DDC
This technical report has been reviewed and is
approved for public release in accordance with AFR 190-12 (7b).
Distribution is unlimited.
A. D. BLOSE
Technical Information Officer

I. INTRODUCTION

The purpose of this correspondence is to extend the results of [1] by considering a more general class of nonlinear control systems. The results obtained in this work provide sufficient conditions for global and local controllability of perturbed nonlinear systems.

Consider the nonlinear time-varying system

$$\frac{d}{dt} x(t) = A(t, x(t), u(t))x(t) + B(t)u(t) + f(t, x(t), u(t)) \quad (1)$$

$t \in [t_0, t_1]$, where $x(t)$ is a $n \times 1$ state vector; $u(t)$ is a $m \times 1$ input vector; A, B, f are $n \times n$, $n \times m$, $n \times 1$ matrix-valued functions, respectively.

Denote $C_{nm}[t_0, t_1]$ as the Banach space of continuous $R^n \times R^m$ valued functions on $[t_0, t_1]$ with the uniform norm $||[x(t), u(t)]|| = |x(t)| + |u(t)|$, where $|w(t)| = \max_i \max_{t \in [t_0, t_1]} |w_i(t)|$ and $|w_i(t)|$ is the absolute value of $w_i(t)$, the element of $w(t)$.

Define the norm of a continuous $n \times m$ matrix-valued function $S(t)$ by $||S(t)|| = \max_i \sum_{j=1}^m \max_{t \in [t_0, t_1]} |S_{ij}(t)|$, where S_{ij} are elements of S .

Given (x_0, x_1) as the initial and final state, respectively of (1). The problem is to find a continuous input function $u(t)$, defined in $[t_0, t_1]$, which steers system (1) from x_0 at t_0 to x_1 at t_1 . The usual definitions of globally and locally completely and totally controllable are assumed [2].

For each fixed element $[z, v] \in C_{nm}[t_0, t_1]$, consider the following system

$$\frac{d}{dt} x(t) = A(t, z(t), v(t))x(t) + B(t)u(t) + f(t, z(t), v(t)) \quad (2)$$

The solution of the system (2) with $x(t_0) = x_0$ is given by

$$\begin{aligned}
 x(t) = & \phi(t, t_0; z, v)x_0 + \int_{t_0}^t \phi(t, s; z, v)B(s)u(s)ds \\
 & + \int_{t_0}^t \phi(t, s; z, v)f(s, z, v)ds
 \end{aligned} \tag{3}$$

where $\phi(t, t_0; z, v)$ is the state transition matrix of the homogeneous system $\frac{d}{dt} x(t) = A(t, z(t), v(t))x(t)$, $\phi(t_0, t_0; z, v) = I$ the identity matrix. Define the controllability matrix by

$$G(t_0, t; z, v) = \int_{t_0}^t \phi(t, s; z, v)B(s)B'(s)\phi'(t, s; z, v)ds \tag{4}$$

the prime denotes the matrix transpose operation. Obviously, $G(t_0, t; z, v)$ is symmetric and non-negative definite.

II. GLOBAL CONTROLLABILITY RESULT

Theorem 1: The system (1) is globally (a) completely controllable at t_0 , or (b) totally controllable if the following conditions are satisfied.

- (i) $B(t)$ has a continuous first derivative with respect to t ,
- (ii) $A(t, x, u)$, $A_x(t, x, u)$, $A_u(t, x, u)$, $f(t, x, u)$, $f_x(t, x, u)$, and $f_u(t, x, u)$ are continuous and bounded in $[t_0, t_1] \times R^n \times R^m$,
- (iii) there exists a positive constant q such that

$$\inf_{[z, v] \in C_{nm}} \det_{[t_0, t_1]} G(t_0, t_1; z, v) \geq q \tag{5}$$

(a) for some $t_1 > t_0$, or (b) for all t_0 and all $t_1 > t_0$.

Proof: The proof of the theorem is based on the Schauder's fixed point theorem of the following version, "Every continuous map which maps a compact convex subset of a Banach space into itself has a fixed point".

For each fixed element $[z, v] \in C_{nm}[t_0, t_1]$, consider the control function $u(t)$ for $t \in [t_0, t_1]$

$$u(t) = B'(t)\phi'(t_1, t; z, v)G^{-1}(t_0, t_1; z, v)[x_1 - \phi(t_1, t_0; z, v)x_0 - \int_{t_0}^{t_1} \phi(t_1, s; z, v)f(s, z, v)ds] \quad (6)$$

where $\phi(t, t_0; z, v)$ is defined as that in Eq. (3). With this expression, Eq. (3) can be rewritten as

$$\begin{aligned} x(t) = & \phi(t, t_0; z, v)x_0 \\ & + \int_{t_0}^t \phi(t, s; z, v)B(s)B'(s)\phi'(t_1, s; z, v)G^{-1}(t_0, t_1; z, v)ds \cdot [x_1 - \phi(t_1, t_0; z, v)x_0 - \\ & - \int_{t_0}^{t_1} \phi(t_1, s; z, v)f(s, z, v)ds] + \int_{t_0}^t \phi(t, s; z, v)f(s, z, v)ds \end{aligned} \quad (7)$$

It should be noted that by hypothesis (iii) $G^{-1}(t_0, t_1; z, v)$ is well-defined in above expression. It is easily seen that $x(t)$ in Eq. (7) satisfies both boundary conditions at $t = t_0$ and $t = t_1$.

Now the right sides of Eqs. (6)-(7) can be viewed as a pair of operators, $P_2([z, v])(t)$ and $P_1([z, v])(t)$, respectively. Define the nonlinear mapping

$$P([z, v])(t) = [P_1([z, v])(t), P_2([z, v])(t)] .$$

Obviously, $P_1([z, v])(t)$ and $P_2([z, v])(t)$ are continuous in t by the uniform continuity of $\phi(t, t_0; z, v)$ in t . Therefore, P maps $C_{nm}[t_0, t_1]$ into $C_{nm}[t_0, t_1]$. It can also be easily verified that, by hypothesis (ii) and the definition of $\phi(t, t_0; z, v)$, P is continuous in $[z, v]$.

Considering the subset of $C_{nm}[t_0, t_1]$

$$S = \left\{ [z, v] \in C_{nm}[t_0, t_1] : ||[z, v]|| \leq K, \right. \\ \left. ||[z(t), v(t)] - [z(\tilde{t}), v(\tilde{t})]|| \leq K|t - \tilde{t}|, \forall t, \tilde{t} \in [t_0, t_1] \right\}$$

where K is certain positive constant depending upon the bounds of A, B, f and their partial derivatives, it can be easily shown that the image set $P(S) \subset S$. Besides, S is closed and convex by this construction. Furthermore, each sequence $\{s_i\}_{i=1}^{\infty} \subset S$ constitutes a uniformly bounded equicontinuous family. Hence by the Arzela-Ascoli theorem [3], S is relative compact and therefore compact.

Then, the Schauder's theorem [3] can be applied to conclude that P has a fixed point $[z^*, v^*]$, i.e.,

$$P([z^*, v^*])(t) = [P_1([z^*, v^*]), P_2([z^*, v^*])] = [z^*, v^*]$$

Substitute this fixed point into Eqs. (6)-(7), a direct differentiation of Eq. (7) with respect to t shows that $z^*(t)$ is a solution to the system (1) for the control function $u(t)$ given by $v^*(t)$.

If condition (iii)(a) holds, $v^*(t)$ drives the system (1) from x_0 to x_1 on some interval $[t_0, t_1]$ for all $x_0, x_1 \in R^n$, system (1) is globally completely controllable at t_0 . If condition (iii)(b) holds, we have global total controllability.

Corollary 1: Given the system (1) with conditions (i)-(iii), then the perturbed system

$$\frac{d}{dt} x(t) = [A(t, x, u) + \epsilon \tilde{A}(t, x, u)]x(t) + [B(t) + \epsilon \tilde{B}(t)]u(t) + f(t, x, u), \quad (8)$$

with \tilde{A}, \tilde{B} satisfying the same type of conditions imposed on A, B , is globally controllable provided ϵ is sufficiently small.

Proof: One needs only to show that the determinant of the modified controllability matrix G^* has a positive infimum. By observing that the determinant of $G^*(t_1, t_0; z, v, \epsilon)$ can be expanded about $\epsilon = 0$ [5] into the sum of $\det G(t_1, t_0; z, v)$, $\epsilon H(t_1, t_0; z, v)$ and $o(\epsilon^2)$ where $H(t_1, t_0; z, v)$ is a bounded scalar, we know that by taking ϵ small enough there exists some $q^* > 0$ such that

$$\det G^*(t_1, t_0; z, v, \epsilon) \geq q^* \quad \text{for all} \quad [z, v] \in C_{nm}[t_0, t_1].$$

Therefore, Theorem 1 concludes the result of controllability.

- Remarks: (i) If (A, f) do not explicitly depend on $u(t)$, the theorem is applicable to the class of systems with $B = B(t, x(t))$. These results then reduce to the work reported in [4].
- (ii) Uniform boundedness of partial derivatives can be slightly weakened by considering a Lipschitz condition on corresponding variables.

III. LOCAL CONTROLLABILITY RESULT

Consider the following subset of $[t_0, t_1] \times R^n \times R^m$,

$$D = \{(t, x, u) : t_0 \leq t \leq t_1, \quad |x| + |u| \leq d, \\ d \text{ is some positive constant}\}.$$

Theorem 2: The system (1) is locally (a) completely controllable at t_0 , or (b) totally controllable if the following conditions are satisfied.

- (i) $B(t)$ has a continuous first derivative with respect to t ,
- (ii) $A(t, x, u)$, $A_x(t, x, u)$, $A_u(t, x, u)$, $f(t, x, u)$, $f_x(t, x, u)$ and $f_u(t, x, u)$ are continuous in D ,

(iii) There exists a positive constant \tilde{q} such that

$$\inf_{[z,v] \in E} \det G(t_1, t_0; z, v) \geq \tilde{q}$$

where

$$E = \{[z,v] \in C_{nm}[t_0, t_1] : ||[z,v]|| \leq d\}$$

and

$$M < \min \left\{ \frac{d}{k_1}, d, \frac{d}{k_2} \right\}$$

where

M is the bound of f in D , k_1 and k_2 are certain positive constants depending upon the bounds of A, B and their partial derivatives in D .

(a) for some $t_1 > t_0$, or (b) for all t_0 and all $t_1 > t_0$.

Proof: The proof is similar to that of Theorem 1. We leave it to the reader.

Corollary 2: Given the system (1) with conditions (i)-(iii) in Theorem 2, the perturbed system

$$\frac{d}{dt} x(t) = [A(t, x, u) + \epsilon \tilde{A}(t, x, u)]x(t) + [B(t) + \epsilon \tilde{B}(t)]u(t) + f(t, x, u) \quad (9)$$

with \tilde{A}, \tilde{B} satisfying the same type of conditions imposed upon A, B is locally controllable provided ϵ is sufficiently small.

Proof: The proof is similar to that of Corollary 1 and follows immediately from Theorem 2.

Example: Consider the system

$$\begin{aligned}\dot{x}_1 &= x_2 + [\sin^2(x_1 + u)]x_2 \\ \dot{x}_2 &= x_1 + u + [\sin^2(x_1 + u)]x_1\end{aligned}$$

In matrix form, we have

$$A(t, x, u) = \begin{bmatrix} 0 & 1 + \sin^2(x_1 + u) \\ 1 + \sin^2(x_1 + u) & 0 \end{bmatrix}, \quad B(t) = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

It is easily seen that $B, dB/dt, A, \partial A/\partial x, \partial A/\partial u$ exist and are bounded in R and $R^2 \times R$, respectively. Furthermore, for each fixed $[z, v] \in C_{21}[t_0, t_1]$, the controllability matrix G is

$$G(t_0, t_1; z, v) = \frac{1}{4} \begin{bmatrix} \int_{t_0}^{t_1} (a-b)^2 dy & \int_{t_0}^{t_1} (a^2 - b^2) dy \\ \int_{t_0}^{t_1} (a^2 - b^2) dy & \int_{t_0}^{t_1} (a+b)^2 dy \end{bmatrix}$$

where

$$a = \exp \int_{t_0}^{t_1} [1 + \sin^2(z_1 + v)] dy, \quad b = \exp \int_{t_0}^{t_1} -[1 + \sin^2(z_1 + v)] dy.$$

It can be easily shown that $\det G(t_0, t_1; z, v) \geq q(t_0, t_1)$ for all $[z, v]$ where $q(t_0, t_1) = \frac{1}{16} [e^{2(t_1 - t_0)} - 1 - 2(t_1 - t_0) - 2(t_1 - t_0)^2 - \frac{4}{3}(t_1 - t_0)^3] > 0$ if $t_1 > t_0$. Hence, by Theorem 1 the system is globally totally controllable.

However, the result of [1] cannot be applied here because in this case the non-linearity $f(t, x, u) = [x_2 \sin^2(x_1 + u), x_1 \sin(x_1 + u)]'$ is not bounded in $R \times R^2 \times R$.

IV. CONCLUSION

Sufficient conditions for global and local controllability of a class of nonlinear systems have been given. A natural question arises as to whether or not these results hold by the same fixed point arguments if the system matrices (A, f) are parameterized as $A(t, p(t), q(t))$ and $f(t, p(t), q(t))$ where $[p, q] \in C_{nm}[t_0, t_1]$ so that for each given $[p, q]$ the parametrized linear time-varying system is controllable. The answer is negative because boundedness and continuity of the partial derivatives of relevant matrices are required in addition to the nonsingularity of the controllability matrix.

Acknowledgment

The author wishes to express his gratitude to Professor A. E. Pearson for his encouragement and helpful comments to this work.

References

- [1] K. Mirza and B. F. Womack, "On the controllability of a class of nonlinear systems", IEEE Trans. Automat. Contr., (Short Paper), Vol. AC-17, pp. 531-535, August 1972.
- [2] E. J. Davison and E. G. Kunze, "Some sufficient conditions for the global and local controllability of nonlinear time-varying systems", SIAM J. Control, Vol. 8, No. 4, pp. 489-497, November 1970.
- [3] L. W. Kantorovich and G. P. Akilov, Functional Analysis in Normed Spaces, New York: McMillan, 1964.
- [4] J. Klamka, "On the controllability of perturbed nonlinear systems", IEEE Trans. Automat. Contr. (Corresp.), Vol. AC-20, pp. 170-172, February 1975.
- [5] E. J. Davison and D. Ball, "On the controllability of perturbed controllable systems", IEEE Trans. Automat. Contr. (Corresp.), Vol. AC-17, pp. 825-826, December 1972.

KCW/llt
3/76

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR-76-1153	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A CLASS OF CONTROLLABLE NONLINEAR SYSTEMS	5. TYPE OF REPORT & PERIOD COVERED Interim rept.	
7. AUTHOR(s) Kuang-Chung Wei	8. CONTRACT OR GRANT NUMBER(s) AFOSR 75-2793	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Brown University Division of Engineering Providence, Rhode Island 02912	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 9769-01	
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Office of Scientific Research (NM) Bolling AFB, DC 20332	12. REPORT DATE 1976	13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) AF-AFOSR-2793-75	15. SECURITY CLASS. (of this Report) UNCLASSIFIED	
15a. DECLASSIFICATION/DOWNGRADING SCHEDULE		
16. DISTRIBUTION STATEMENT (of this Report) 9769 17 01		
Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Sufficient conditions for global and local controllability of the nonlinear system $\frac{d}{dt}x(t) = A(t, x(t), u(t))x(t) + B(t)u(t) + f(t, x(t), u(t))$ as its perturbed system are given. These conditions extend some previous results through the removal of certain boundedness conditions involving the functions (A, f) and their partial derivatives.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)